

מרץ 2022

רופא/ה יקר/ה
חוקח/ת יקר/ה,**KEYTRUDA® 100 mg/4 mL**
קיטרודה 100 מ"ג/א"מ 4 מ"ל

Dosage form and Composition:

Pembrolizumab 100 mg/4 ml; Concentrate for Solution for Intravenous Infusion

חברת מרק שארפ ודוחם (ישראל-1996) בע"מ, מבקשת לידע על עדכון העלון לרופא של Keytruda 100mg/4ml

עדכנים מוחשיים שבעצם לעלון לרופא (טקסט שהוסף לעלון לרופא מודגש בקו תחתון, טקסט שנמחק מהעלון לרופא מסומן בקו חוצה):

1 THERAPEUTIC INDICATIONS

1.16 Cutaneous Squamous Cell Carcinoma

KEYTRUDA is indicated for the treatment of patients with recurrent or metastatic cutaneous squamous cell carcinoma (cSCC) or locally advanced cSCC that is not curable by surgery or radiation.

6 ADVERSE REACTIONS

6.1 Clinical Trials Experience

העדכנים בפרק זה כוללים עדכון מידע לגבי התוויות cSCC.

14 CLINICAL STUDIES

עדכון מידע לגבי התוויות:

Microsatellite Instability-High or Mismatch Repair Deficient Colorectal Cancer (CRC)
Cutaneous Squamous Cell Carcinoma

התוויות המאושרות לתקשייה:

Melanoma

- KEYTRUDA (pembrolizumab) is indicated for the treatment of patients with unresectable or metastatic melanoma.
- KEYTRUDA is indicated for the adjuvant treatment of patients with melanoma with involvement of lymph node (s) following complete resection.

Non-Small Cell Lung Cancer

- KEYTRUDA, in combination with pemetrexed and carboplatin, is indicated for the first-line treatment of patients with metastatic nonsquamous non-small cell lung cancer (NSCLC) negative for EGFR or ALK genomic tumor aberrations.
- KEYTRUDA, in combination with carboplatin and either paclitaxel or paclitaxel protein-bound, is indicated for the first-line treatment of patients with metastatic squamous NSCLC.
- KEYTRUDA, as a single agent, is indicated for the treatment of patients with metastatic NSCLC whose tumors express PD-L1 [Tumor Proportion Score (TPS) $\geq 50\%$] as determined by a validated test. Patients with EGFR or ALK genomic tumor aberrations should have disease progression on or after platinum-containing chemotherapy and an approved therapy for these aberrations prior to receiving KEYTRUDA.
- KEYTRUDA, as a single agent, is indicated for the treatment of patients with advanced NSCLC whose tumors express PD-L1 as determined by a validated test, with disease progression on or after platinum containing chemotherapy. Patients with EGFR or ALK genomic tumor aberrations should have disease progression on approved therapy for these aberrations prior to receiving KEYTRUDA [see Clinical Studies (14.2)].

Small Cell Lung Cancer

KEYTRUDA is indicated for the treatment of patients with metastatic small cell lung cancer (SCLC) with disease progression on or after platinum-based chemotherapy and at least one other prior line of therapy, that have not been previously treated with immunotherapy.

Head and Neck Cancer

- KEYTRUDA, in combination with platinum and fluorouracil (FU), is indicated for the first-line treatment of patients with metastatic or with unresectable, recurrent head and neck squamous cell carcinoma (HNSCC).
- KEYTRUDA, as a single agent, is indicated for the first-line treatment of patients with metastatic or with unresectable, recurrent HNSCC whose tumors express PD-L1 [Combined Positive Score (CPS) ≥ 1] as determined by a validated test.
- KEYTRUDA is indicated for the treatment of patients with recurrent or metastatic head and neck squamous cell carcinoma (HNSCC) with disease progression on or after platinum-containing chemotherapy.

Classical Hodgkin Lymphoma

KEYTRUDA is indicated for the treatment of adult patients with relapsed or refractory classical Hodgkin lymphoma (cHL).

KEYTRUDA is indicated for the treatment of pediatric patients with refractory cHL, or cHL that has relapsed after 2 or more lines of therapy.

Primary Mediastinal large B-Cell Lymphoma

KEYTRUDA is indicated for the treatment of adult and pediatric patients with refractory primary mediastinal large B-cell lymphoma (PMBCL), or who have relapsed after 2 or more prior lines of therapy.

Limitation of Use: KEYTRUDA is not recommended for treatment of patients with PMBCL who require urgent cytoreductive therapy.

Urothelial Carcinoma

•KEYTRUDA is indicated for the treatment of patients with locally advanced or metastatic urothelial carcinoma who are not eligible for cisplatin-containing chemotherapy and whose tumors express PD-L1 [Combined Positive Score (CPS ≥ 10)] as determined by a validated test, or in patients who are not eligible for any platinum-containing chemotherapy regardless of PD-L1 status.

•KEYTRUDA is indicated for the treatment of patients with locally advanced or metastatic urothelial carcinoma who have disease progression during or following platinum-containing chemotherapy or within 12 months of neoadjuvant or adjuvant treatment with platinum-containing chemotherapy.

Non-Muscle Invasive Bladder Cancer (NMIBC)

KEYTRUDA is indicated for the treatment of patients with Bacillus Calmette-Guerin (BCG)-unresponsive, high-risk, non-muscle invasive bladder cancer (NMIBC) with carcinoma in situ (CIS) with or without papillary tumors who are ineligible for or have elected not to undergo cystectomy.

Microsatellite Instability-High Cancer

KEYTRUDA is indicated for the treatment of adult and pediatric patients with unresectable or metastatic, microsatellite instability-high (MSI H) or mismatch repair deficient (dMMR).

- solid tumors that have progressed following prior systemic treatment and who have no satisfactory alternative treatment options, or
- colorectal cancer that has progressed following treatment with a fluoropyrimidine, oxaliplatin, and irinotecan.

Limitation of Use: The safety and effectiveness of KEYTRUDA in pediatric patients with MSI H central nervous system cancers have not been established.

Gastric Cancer

KEYTRUDA is indicated for the treatment of patients with recurrent locally advanced or metastatic gastric or gastroesophageal junction adenocarcinoma whose tumors express PD-L1 [Combined Positive Score (CPS) ≥ 1] as determined by a validated test, with disease progression on or after two or more prior lines of therapy including fluoropyrimidine- and platinum-containing chemotherapy and if appropriate, HER2/neu targeted therapy.

Cervical Cancer

KEYTRUDA is indicated for the treatment of patients with recurrent or metastatic cervical cancer with disease progression on or after chemotherapy whose tumors express PD-L1 (CPS ≥ 1) as determined by a validated test.

Merkel Cell Carcinoma

KEYTRUDA is indicated for the treatment of adult and pediatric patients with recurrent locally advanced or metastatic Merkel cell carcinoma (MCC).

Renal Cell Carcinoma

KEYTRUDA, in combination with axitinib, is indicated for the first-line treatment of patients with advanced renal cell carcinoma (RCC).



KEYTRUDA, in combination with lenvatinib, is indicated for the first-line treatment of adult patients with advanced RCC.

Tumor Mutational Burden-High (TMB-H) Cancer

KEYTRUDA is indicated for the treatment of adult and pediatric patients with unresectable or metastatic tumor mutational burden-high (TMB-H) [≥ 10 mutations/megabase (mut/Mb)] solid tumors, as determined by a validated test, that have progressed following prior treatment and who have no satisfactory alternative treatment options.

Limitations of Use: The safety and effectiveness of KEYTRUDA in pediatric patients with TMB-H central nervous system cancers have not been established.

Esophageal Cancer

•KEYTRUDA is indicated for the treatment of patients with locally advanced or metastatic esophageal or gastroesophageal junction (GEJ) (Siewert type I) carcinoma that is not amenable to surgical resection or definitive chemoradiation in combination with platinum- and fluoropyrimidine-based chemotherapy.

•KEYTRUDA is indicated for the treatment of patients with recurrent locally advanced or metastatic squamous cell carcinoma of the esophagus whose tumors express PD-L1 (CPS ≥ 10) as determined by a validated test, with disease progression after one or more prior lines of systemic therapy.

Cutaneous Squamous Cell Carcinoma

KEYTRUDA is indicated for the treatment of patients with recurrent or metastatic cutaneous squamous cell carcinoma (cSCC) or locally advanced cSCC that is not curable by surgery or radiation.

Microsatellite Instability-High or Mismatch Repair Deficient Colorectal Cancer (CRC)

KEYTRUDA is indicated for the first-line treatment of patients with unresectable or metastatic MSI-H or dMMR colorectal cancer (CRC).

Triple negative breast cancer (TNBC)

KEYTRUDA, in combination with chemotherapy, is indicated for the treatment of patients with locally recurrent unresectable or metastatic triple negative breast cancer (TNBC) whose tumors express PD-L1 (CPS ≥ 10) as determined by a validated test.

Endometrial carcinoma

Keytruda, in combination with lenvatinib, is indicated for the treatment of advanced or recurrent endometrial carcinoma in adults who have disease progression on or following prior treatment with a platinum containing therapy and who are not candidates for curative surgery or radiation.

ל מידע מלא ולהוראות מתן מפורטות, יש לעיין בעלון לרופא המאושר על ידי משרד הבריאות.

העלון לרופא ולצורך נשלח לפרסום במאגר התרופות שבמשרד הבריאות, וניתן לקבלו מודפס על ידי פניה לעל הרישום, חברת MSD, טלפון .09-9533333 מופצת ע"י חברת נובלוג בע"מ.

בברכה,
דורות מאורי
חזקת ממוני
ישראל MSD

References:

Keytruda_100mg_4ml-SPC-03_2022